

SEATTLE SPERM BANK

Attn: Dr. Jeffrey Olliffe 4915 25th Ave NE, Suite 204W

Seattle, WA 98105 Phone: (206) 588-1484 Fax: (206) 466-4696 NPI: 1306838271 Report Date: 12/18/2017 MALE

DONOR 10219

•

Ethnicity: Mixed or Other

Caucasian

Sample Type: EDTA Blood Date of Collection: 12/08/2017 Date Received: 12/11/2017 Date Tested: 12/18/2017 Barcode: 11004212319195 Accession ID: CSL4XXP23CJZV2K Indication: Egg or sperm donor FEMALE N/A

Foresight™ Carrier Screen

NEGATIVE

ABOUT THIS TEST

The **Counsyl Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

RESULTS SUMMARY

Risk Details	DONOR 10219	Partner
Panel Information	Foresight Carrier Screen Universal Panel Minus X-Linked (102 conditions tested)	N/A
All conditions tested A complete list of all conditions tested can be found on page 4.	 □ NEGATIVE No disease-causing mutations were detected. 	N/A

CLINICAL NOTES NEXT STEPS

None

• If necessary, patients can discuss residual risks with their physician or a genetic counselor.

Counsyl has renamed its products effective July 19, 2017. The Family Prep Screen is now the Foresight Carrier Screen. The new names now appear on all communications from Counsyl. If you have any questions, please contact Counsyl directly.



RESULTS RECIPIENT

SEATTLE SPERM BANK

Attn: Dr. Jeffrey Olliffe

NPI: 1306838271

Report Date: 12/18/2017

MALE
DONOR 10219
DOB:

Ethnicity: Mixed or Other

Caucasian

Barcode: 11004212319195

FEMALE N/A

Methods and Limitations

DONOR 10219 [Foresight Carrier Screen]: sequencing with copy number analysis, spinal muscular atrophy, and analysis of homologous regions.

Sequencing with copy number analysis

High-throughput sequencing and read depth-based copy number analysis are used to analyze the listed exons, as well as selected intergenic and intronic regions, of the genes in the Conditions Tested section of the report. The region of interest (ROI) of the test comprises these regions, in addition to the 20 intronic bases flanking each exon. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to high coverage and the sequences are compared to standards and references of normal variation. More than 99% of all bases in the ROI are sequenced at greater than the minimum read depth. Mutations may not be detected in areas of lower sequence coverage. Small insertions and deletions may not be as accurately determined as single nucleotide variants. Genes that have closely related pseudogenes may be addressed by a different method. *CFTR* and *DMD* testing includes analysis for both large (exon-level) deletions and duplications with an average sensitivity of 99%, while other genes are only analyzed for large deletions with a sensitivity of >75%. However, the sensitivity may be higher for selected founder deletions. If *GJB2* is tested, two large upstream deletions which overlap *GJB6* and affect the expression of *GJB2*, del(*GJB6*-D13S1830) and del(*GJB6*-D13S1854), are also analyzed. Mosaicism or somatic variants present at low levels may not be detected. If detected, these may not be reported.

Detection rates are determined by using literature to estimate the fraction of disease alleles, weighted by frequency, that the methodology is unable to detect. Detection rates only account for analytical sensitivity and certain variants that have been previously described in the literature may not be reported if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease-specific rates of de novo mutations.

All variants that are a recognized cause of the disease will be reported. In addition, variants that have not previously been established as a recognized cause of disease may be identified. In these cases, only variants classified as "likely" pathogenic are reported. Likely pathogenic variants are described elsewhere in the report as "likely to have a negative impact on gene function". Likely pathogenic variants are evaluated and classified by assessing the nature of the variant and reviewing reports of allele frequencies in cases and controls, functional studies, variant annotation and effect prediction, and segregation studies. Exon level duplications are assumed to be in tandem and are classified according to their predicted effect on the reading frame. Benign variants, variants of uncertain significance, and variants not directly associated with the intended disease phenotype are not reported. Curation summaries of reported variants are available upon request.

Spinal muscular atrophy

Targeted copy number analysis is used to determine the copy number of exon 7 of the *SMN1* gene relative to other genes. Other mutations may interfere with this analysis. Some individuals with two copies of *SMN1* are carriers with two *SMN1* genes on one chromosome and a *SMN1* deletion on the other chromosome. This is more likely in individuals who have 2 copies of the *SMN1* gene and are positive for the g.27134T>G SNP, which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have 2 copies of *SMN1*.

Analysis of homologous regions

A combination of high-throughput sequencing, read depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss of function mutations in certain genes that have homology to other regions. The precise breakpoints of large deletions in these genes cannot be determined, but are estimated from copy number analysis. High numbers of pseudogene copies may interfere with this analysis.

If *CYP21A2* is tested, patients who have one or more additional copies of the *CYP21A2* gene and a loss of function mutation may not actually be a carrier of 21-hydroxylase-deficient congenital adrenal hyperplasia (CAH). Because the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are only based on published incidences for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate of overall chances for 21-hydroxylase-deficient CAH, especially in the aforementioned populations, as they do not account for non-classic CAH. If *HBA11HBA2* are tested, some individuals with four alpha globin genes may be carriers, with three genes on one chromosome and a deletion on the other chromosome. This and similar, but rare, carrier states, where complementary changes exist in both the gene and a pseudogene, may not be detected by the assay.



RESULTS RECIPIENT **SEATTLE SPERM BANK** Attn: Dr. Jeffrey Olliffe NPI: 1306838271

Report Date: 12/18/2017

MALE **DONOR 10219** DOB

Ethnicity: Mixed or Other Caucasian

Barcode: 11004212319195

FEMALE N/A

Limitations

In an unknown number of cases, nearby genetic variants may interfere with mutation detection. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions and technical errors. This test is designed to detect and report germline alterations. While somatic variants present at low levels may be detected, these may not be reported. If more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes. The test does not fully address all inherited forms of intellectual disability, birth defects and genetic disease. A family history of any of these conditions may warrant additional evaluation. Furthermore, not all mutations will be identified in the genes analyzed and additional testing may be beneficial for some patients. For example, individuals of African, Southeast Asian, and Mediterranean ancestry are at increased risk for being carriers for hemoglobinopathies, which can be identified by CBC and hemoglobin electrophoresis or HPLC (ACOG Practice Bulletin No. 78. Obstet. Gynecol. 2007;109:229-37).

This test was developed and its performance characteristics determined by Counsyl, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: #05D1102604.

LAB DIRECTORS

H. Peter Kang, MD, MS, FCAP

Hyunseok Kang



SEATTLE SPERM BANK Attn: Dr. Jeffrey Olliffe

NPI: 1306838271 Report Date: 12/18/2017 **DONOR 10219**

MALE

DOB:

Ethnicity: Mixed or Other

Caucasian

Barcode: 11004212319195

FEMALE N/A

Conditions Tested

21-hydroxylase-deficient Congenital Adrenal Hyperplasia - Gene: CYP21A2. Autosomal Recessive. Analysis of Homologous Regions. Variants (13): CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs*21, I173N, L308FfsX6, P31L, Q319*, Q319*+CYP21A2dup, R357W, V281L, [I237N;V238E;M240K], c.293-13C>G. Detection Rate: Mixed or Other Caucasian 96%.

ABCC8-related Hyperinsulinism - Gene: ABCC8. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000352:1-39. Detection Rate: Mixed or Other Caucasian >99%.

Alkaptonuria - Gene: HGD. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000187:1-14. Detection Rate: Mixed or Other Caucasian >99%. Alpha Thalassemia - Genes: HBA1, HBA2. Autosomal Recessive. Analysis of Homologous Regions. Variants (13): -(alpha)20.5, --BRIT, --MEDI, --MEDI, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, anti3.7, anti4.2. del HS-40. Detection Rate: Unknown due to rarity of disease.

Alpha-1 Antitrypsin Deficiency - Gene: SERPINA1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000295:2-5. Detection Rate: Mixed or Other Caucasian >99%.

Alpha-mannosidosis - Gene: MAN2B1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000528:1-23. Detection Rate: Mixed or Other Caucasian >99%

Alpha-sarcoglycanopathy - Gene: SGCA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000023:1-9. Detection Rate: Mixed or Other

Andermann Syndrome - Gene: SLC12A6. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_133647:1-25. Detection Rate: Mixed or Other Caucasian >99%.

ARSACS - Gene: SACS. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_014363:2-10. Detection Rate: Mixed or Other Caucasian 99%. Aspartylglycosaminuria - Gene: AGA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000027:1-9. Detection Rate: Mixed or Other Caucasian >99%.

Ataxia with Vitamin E Deficiency - Gene: TTPA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000370:1-5. Detection Rate: Mixed or Other Caucasian >99%.

Ataxia-telangiectasia - Gene: ATM. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000051:2-63. Detection Rate: Mixed or Other

Bardet-Biedl Syndrome, BBS1-related - Gene: BBS1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_024649:1-17. Detection Rate: Mixed or Other Caucasian >99%.

Bardet-Biedl Syndrome, BBS10-related - Gene: BBS10. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_024685:1-2. Detection Rate: Mixed or Other Caucasian >99%.

Beta-sarcoglycanopathy - Gene: SGCB. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000232:1-6. Detection Rate: Mixed or Other Caucasian >99%

Biotinidase Deficiency - Gene: BTD. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000060:1-4. Detection Rate: Mixed or Other Caucasian >99%.

Bloom Syndrome - Gene: BLM. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000057:2-22. Detection Rate: Mixed or Other

Canavan Disease - Gene: ASPA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000049:1-6. Detection Rate: Mixed or Other Caucasian 98%.

Carnitine Palmitoyltransferase IA Deficiency - Gene: CPT1A. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_001876:2-19. Detection Rate: Mixed or Other Caucasian >99%.

Carnitine Palmitoyltransferase II Deficiency - Gene: CPT2. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000098:1-5. Detection Rate: Mixed or Other Caucasian >99%.

Cartilage-hair Hypoplasia - Gene: RMRP. Autosomal Recessive. Sequencing with Copy Number Analysis. Exon: NR_003051:1. Detection Rate: Mixed or Other Caucasian >99%.

Citrullinemia Type 1 - Gene: ASS1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000050:3-16. Detection Rate: Mixed or Other Caucasian >99%.

CLN3-related Neuronal Ceroid Lipofuscinosis - Gene: CLN3. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_001042432:2-16. Detection Rate: Mixed or Other Caucasian >99%.

CLN5-related Neuronal Ceroid Lipofuscinosis - Gene: CLN5. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_006493:1-4. Detection Rate: Mixed or Other Caucasian >99%.

CNGB3-related Achromatopsia - Gene: CNGB3. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_019098:1-18. Detection Rate: Mixed or Other Caucasian >99%.

Cohen Syndrome - Gene: VPS13B. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_017890:2-62. Detection Rate: Mixed or Other Caucasian 97%.

Congenital Disorder of Glycosylation Type Ia - Gene: PMM2. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000303:1-8. Detection Rate: Mixed or Other Caucasian >99%.

Congenital Disorder of Glycosylation Type Ib - Gene: MPI. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_002435:1-8. Detection Rate: Mixed or Other Caucasian >99%.

Congenital Finnish Nephrosis - Gene: NPHS1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_004646:1-29. Detection Rate: Mixed or Other Caucasian >99%.

Costeff Optic Atrophy Syndrome - Gene: OPA3. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_025136:1-2. Detection Rate: Mixed or Other Caucasian >99%.

Cystic Fibrosis - Gene: CFTR. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. Detection Rate: Mixed or Other Caucasian >99% Cystinosis - Gene: CTNS. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_004937:3-12. Detection Rate: Mixed or Other Caucasian >99%. D-bifunctional Protein Deficiency - Gene: HSD17B4. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000414:1-24. Detection Rate: Mixed or Other Caucasian 98%.

Dihydropyrimidine Dehydrogenase Deficiency - Gene: DPYD. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000110:1-23. Detection Rate: Mixed or Other Caucasian 98%.

Factor XI Deficiency - Gene: F11. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000128:2-15. Detection Rate: Mixed or Other Caucasian >99%

Familial Dysautonomia - Gene: IKBKAP. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_003640:2-37. Detection Rate: Mixed or Other Caucasian >99%.

Familial Mediterranean Fever - Gene: MEFV. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000243:1-10. Detection Rate: Mixed or

Fanconi Anemia Type C - Gene: FANCC. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000136:2-15. Detection Rate: Mixed or Other Caucasian >99%

FKTN-related Disorders - Gene: FKTN. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_001079802:3-11. Detection Rate: Mixed or Other Caucasian >99%.

Galactosemia - Gene: GALT. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000155:1-11. Detection Rate: Mixed or Other Caucasian >99%. Gaucher Disease - Gene: GBA. Autosomal Recessive. Analysis of Homologous

Regions. Variants (10): D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs*18. Detection Rate: Mixed or Other Caucasian 60%.

GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness - Gene: GJB2. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_004004:1-2. Detection Rate: Mixed or Other Caucasian >99%.

Glutaric Acidemia Type 1 - Gene: GCDH. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000159:2-12. Detection Rate: Mixed or Other Caucasian >99%.



SEATTLE SPERM BANK Attn: Dr. Jeffrey Olliffe

NPI: 1306838271 Report Date: 12/18/2017

MALE

Caucasian

DONOR 10219

Barcode: 11004212319195

DOB: Ethnicity: Mixed or Other

Glycogen Storage Disease Type Ia - Gene: G6PC. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000151:1-5. Detection Rate: Mixed or Other Caucasian >99%.

Glycogen Storage Disease Type Ib - Gene: SLC37A4. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_001164277:3-11. Detection Rate: Mixed or Other Caucasian >99%.

Glycogen Storage Disease Type III - Gene: AGL. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000642:2-34. Detection Rate: Mixed or

Glycogen Storage Disease Type V - Gene: PYGM. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_005609:1-20. Detection Rate: Mixed or Other Caucasian >99%.

GRACILE Syndrome - Gene: BCS1L. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_004328:3-9. Detection Rate: Mixed or Other Caucasian >99%

HADHA-related Disorders - Gene: HADHA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000182:1-20. Detection Rate: Mixed or Other Caucasian >99%

Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease) - Gene: HBB. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000518:1-3. Detection Rate: Mixed or Other Caucasian >99%

Hereditary Fructose Intolerance - Gene: ALDOB. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000035:2-9. Detection Rate: Mixed or Other Caucasian >99%.

Herlitz Junctional Epidermolysis Bullosa, LAMA3-related - Gene: LAMA3. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000227:1-38. Detection Rate: Mixed or Other Caucasian >99%

Herlitz Junctional Epidermolysis Bullosa, LAMB3-related - Gene: LAMB3. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000228:2-23. Detection Rate: Mixed or Other Caucasian >99%.

Herlitz Junctional Epidermolysis Bullosa, LAMC2-related - Gene: LAMC2.

Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM 005562:1-23. Detection Rate: Mixed or Other Caucasian >99%.

Hexosaminidase A Deficiency (Including Tay-Sachs Disease) - Gene: HEXA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000520:1-14. Detection Rate: Mixed or Other Caucasian >99%.

Homocystinuria Caused by Cystathionine Beta-synthase Deficiency - Gene: CBS. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000071:3-17. Detection Rate: Mixed or Other Caucasian >99%

Hypophosphatasia, Autosomal Recessive - Gene: ALPL. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000478:2-12. Detection Rate: Mixed or Other Caucasian >99%.

Inclusion Body Myopathy 2 - Gene: GNE. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_001128227:1-12. Detection Rate: Mixed or Other Caucasian >99%.

Isovaleric Acidemia - Gene: IVD. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_002225:1-12. Detection Rate: Mixed or Other Caucasian >99%

Joubert Syndrome 2 - Gene: TMEM216. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_001173990:1-5. Detection Rate: Mixed or Other Caucasian >99%

Krabbe Disease - Gene: GALC. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000153:1-17. Detection Rate: Mixed or Other Caucasian >99%. Lipoamide Dehydrogenase Deficiency - Gene: DLD. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000108:1-14. Detection Rate: Mixed or Other Caucasian >99%.

Maple Syrup Urine Disease Type 1B - Gene: BCKDHB. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_183050:1-10. Detection Rate: Mixed or Other Caucasian >99%

Medium Chain Acyl-CoA Dehydrogenase Deficiency - Gene: ACADM. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000016:1-12. Detection Rate: Mixed or Other Caucasian >99%.

Megalencephalic Leukoencephalopathy with Subcortical Cysts - Gene: MLC1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons:

NM_015166:2-12. Detection Rate: Mixed or Other Caucasian >99%.

Metachromatic Leukodystrophy - Gene: ARSA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000487:1-8. Detection Rate: Mixed or Other Caucasian >99%.

Mucolipidosis IV - Gene: MCOLN1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_020533:1-14. Detection Rate: Mixed or Other Caucasian >99%.

FEMALE

N/A

Mucopolysaccharidosis Type I - Gene: IDUA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000203:1-14. Detection Rate: Mixed or Other Caucasian >99%.

Muscle-eye-brain Disease - Gene: POMGNT1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_017739:2-22. Detection Rate: Mixed or Other Caucasian 96%.

NEB-related Nemaline Myopathy - Gene: NEB. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_001271208:3-80,117-183. Detection Rate: Mixed or Other Caucasian 92%.

Niemann-Pick Disease Type C - Gene: NPC1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000271:1-25. Detection Rate: Mixed or Other Caucasian >99%.

Niemann-Pick Disease, SMPD1-associated - Gene: SMPD1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000543:1-6. Detection Rate: Mixed or Other Caucasian >99%.

Nijmegen Breakage Syndrome - Gene: NBN. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_002485:1-16. Detection Rate: Mixed or Other Caucasian >99%.

Northern Epilepsy - Gene: CLN8. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_018941:2-3. Detection Rate: Mixed or Other Caucasian >99%.

PCDH15-related Disorders - Gene: PCDH15. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_033056:2-33. Detection Rate: Mixed or Other Caucasian 93%.

Pendred Syndrome - Gene: SLC26A4. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000441:2-21. Detection Rate: Mixed or Other Caucasian >99%.

PEX1-related Zellweger Syndrome Spectrum - Gene: PEX1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000466:1-24. Detection Rate: Mixed or Other Caucasian >99%.

Phenylalanine Hydroxylase Deficiency - Gene: PAH. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000277:1-13. Detection Rate: Mixed or Other Caucasian >99%

PKHD1-related Autosomal Recessive Polycystic Kidney Disease - Gene: PKHD1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM 138694:2-67. Detection Rate: Mixed or Other Caucasian >99%.

Polyglandular Autoimmune Syndrome Type 1 - Gene: AIRE. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000383:1-14. Detection Rate: Mixed or Other Caucasian >99%.

Pompe Disease - Gene: GAA. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM 000152:2-20. Detection Rate: Mixed or Other Caucasian 98%. PPT1-related Neuronal Ceroid Lipofuscinosis - Gene: PPT1. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000310:1-9. Detection Rate: Mixed or Other Caucasian >99%.

Primary Carnitine Deficiency - Gene: SLC22A5. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_003060:1-10. Detection Rate: Mixed or Other Caucasian >99%.

Primary Hyperoxaluria Type 1 - Gene: AGXT. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000030:1-11. Detection Rate: Mixed or Other Caucasian >99%.

Primary Hyperoxaluria Type 2 - Gene: GRHPR. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_012203:1-9. Detection Rate: Mixed or Other Caucasian >99%.

PROP1-related Combined Pituitary Hormone Deficiency - Gene: PROP1.

Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_006261:1-3. Detection Rate: Mixed or Other Caucasian >99%

Pseudocholinesterase Deficiency - Gene: BCHE. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000055:2-4. Detection Rate: Mixed or Other Caucasian >99%.

Pycnodysostosis - Gene: CTSK. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000396:2-8. Detection Rate: Mixed or Other

Rhizomelic Chondrodysplasia Punctata Type 1 - Gene: PEX7. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000288:1-10. Detection Rate: Mixed or Other Caucasian >99%.



SEATTLE SPERM BANK Attn: Dr. Jeffrey Olliffe **NPI:** 1306838271

Report Date: 12/18/2017

MALE
DONOR 10219

DOB: Ethnicity: Mixed or Other

Caucasian

Barcode: 11004212319195

FEMALE N/A

Salla Disease - **Gene:** SLC17A5. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons:** NM_012434:1-11. **Detection Rate:** Mixed or Other Caucasian 98%.

Segawa Syndrome - **Gene**: TH. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons**: NM_000360:1-13. **Detection Rate**: Mixed or Other Caucasian >99%.

Short Chain Acyl-CoA Dehydrogenase Deficiency - Gene: ACADS. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000017:1-10. Detection Rate: Mixed or Other Caucasian >99%.

Sjogren-Larsson Syndrome - **Gene**: ALDH3A2. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons**: NM_000382:1-10. **Detection Rate**: Mixed or Other Caucasian 97%.

Smith-Lemli-Opitz Syndrome - **Gene:** DHCR7. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons:** NM_001360:3-9. **Detection Rate:** Mixed or Other Caucasian >99%.

Spinal Muscular Atrophy - Gene: SMN1. Autosomal Recessive. Spinal Muscular Atrophy. Variant (1): SMN1 copy number. Detection Rate: Mixed or Other Caucasian 95%

Steroid-resistant Nephrotic Syndrome - **Gene:** NPHS2. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons:** NM_014625:1-8. **Detection Rate:** Mixed or Other Caucasian >99%.

Sulfate Transporter-related Osteochondrodysplasia - Gene: SLC26A2. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000112:2-3. Detection Rate: Mixed or Other Caucasian >99%.

TPP1-related Neuronal Ceroid Lipofuscinosis - Gene: TPP1. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons**: NM_000391:1-13. **Detection Rate**: Mixed or Other Caucasian >99%.

Tyrosinemia Type I - Gene: FAH. Autosomal Recessive. Sequencing with Copy Number Analysis. Exons: NM_000137:1-14. Detection Rate: Mixed or Other Caucasian >99%.

Usher Syndrome Type 3 - **Gene:** CLRN1. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons:** NM_174878:1-3. **Detection Rate:** Mixed or Other Caucasian >99%.

Very Long Chain Acyl-CoA Dehydrogenase Deficiency - **Gene**: ACADVL. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons**: NM_000018:1-20. **Detection Rate**: Mixed or Other Caucasian >99%.

Wilson Disease - **Gene**: ATP7B. Autosomal Recessive. Sequencing with Copy Number Analysis. **Exons**: NM_000053:1-21. **Detection Rate**: Mixed or Other Caucasian >99%.



RESULTS RECIPIENT

SEATTLE SPERM BANK

Attn: Dr. Jeffrey Olliffe

NPI: 1306838271

Report Date: 12/18/2017

MALE DONOR 10219

DOB: Ethnicity: Mixed or Other

Caucasian

Barcode: 11004212319195

FEMALE N/A

Risk Calculations

Below are the risk calculations for all conditions tested. Since negative results do not completely rule out the possibility of being a carrier, the **residual risk** represents the patient's post-test likelihood of being a carrier and the **reproductive risk** represents the likelihood the patient's future children could inherit each disease. These risks are inherent to all carrier screening tests, may vary by ethnicity, are predicated on a negative family history and are present even after a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

Disease	DONOR 10219 Residual Risk	Reproductive Risk
21-hydroxylase-deficient Congenital Adrenal Hyperplasia	1 in 1,400	1 in 310,000
ABCC8-related Hyperinsulinism	1 in 11,000	< 1 in 1,000,000
Alkaptonuria	1 in 6,800	< 1 in 1,000,000
lpha Thalassemia	Alpha globin status: aa/aa.	Not calculated
lpha-1 Antitrypsin Deficiency	1 in 2,700	1 in 300,000
lpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Ipha-sarcoglycanopathy	1 in 45,000	< 1 in 1,000,000
ndermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
RSACS	< 1 in 44,000	< 1 in 1,000,000
spartylglycosaminuria	< 1 in 50,000	< 1 in 1,000,000
staxia with Vitamin E Deficiency	< 1 in 50,000	< 1 in 1,000,000
taxia-telangiectasia	1 in 8,200	< 1 in 1,000,000
ardet-Biedl Syndrome, BBS1-related	1 in 16,000	< 1 in 1,000,000
ardet-Biedl Syndrome, BBS10-related	1 in 16,000	< 1 in 1,000,000
eta-sarcoglycanopathy	< 1 in 50,000	< 1 in 1,000,000
iotinidase Deficiency	1 in 13,000	1 in 670,000
loom Syndrome	< 1 in 50,000	< 1 in 1,000,000
anavan Disease	< 1 in 31,000	< 1 in 1,000,000 < 1 in 1,000,000
anavan Disease arnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	
•		< 1 in 1,000,000
arnitine Palmitoyltransferase II Deficiency	< 1 in 50,000	< 1 in 1,000,000
artilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
trullinemia Type 1	1 in 12,000	< 1 in 1,000,000
LN3-related Neuronal Ceroid Lipofuscinosis	1 in 22,000	< 1 in 1,000,000
LN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
NGB3-related Achromatopsia	1 in 11,000	< 1 in 1,000,000
ohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
ongenital Disorder of Glycosylation Type Ia	1 in 16,000	< 1 in 1,000,000
ongenital Disorder of Glycosylation Type Ib	< 1 in 50,000	< 1 in 1,000,000
ongenital Finnish Nephrosis	< 1 in 50,000	< 1 in 1,000,000
osteff Optic Atrophy Syndrome	< 1 in 50,000	< 1 in 1,000,000
ystic Fibrosis	1 in 2,700	1 in 290,000
ystinosis	1 in 22,000	< 1 in 1,000,000
-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000
ihydropyrimidine Dehydrogenase Deficiency	< 1 in 29,000	< 1 in 1,000,000
actor XI Deficiency	< 1 in 50,000	< 1 in 1,000,000
amilial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
amilial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
anconi Anemia Type C	1 in 16,000	< 1 in 1,000,000
KTN-related Disorders	< 1 in 50,000	< 1 in 1,000,000
alactosemia	1 in 8,600	< 1 in 1,000,000
aucher Disease	1 in 280	1 in 120,000
JB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 3,200	1 in 420,000
lutaric Acidemia Type 1	1 in 10,000	< 1 in 1,000,000
•	1 in 18,000	
lycogen Storage Disease Type Ia lycogen Storage Disease Type Ib	1 in 18,000 1 in 35,000	< 1 in 1,000,000 < 1 in 1,000,000
, ,	· · · · · · · · · · · · · · · · · · ·	
ilycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
ilycogen Storage Disease Type V	1 in 16,000	< 1 in 1,000,000
GRACILE Syndrome	< 1 in 50,000	< 1 in 1,000,000
IADHA-related Disorders	1 in 15,000	< 1 in 1,000,000



RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Dr. Jeffrey Olliffe
NPI: 1306838271
Report Date: 12/18/2017

MALE
DONOR 10219
DOB:

Ethnicity: Mixed or Other

Caucasian

Barcode: 11004212319195

FEMALE N/A

Disease	DONOR 10219 Residual Risk	Reproductive Risk
lb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and iickle Cell Disease)	1 in 5,000	1 in 990,000
lereditary Fructose Intolerance	1 in 8,000	< 1 in 1,000,000
erlitz Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
erlitz Junctional Epidermolysis Bullosa, LAMB3-related	< 1 in 50,000	< 1 in 1,000,000
erlitz Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
exosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
omocystinuria Caused by Cystathionine Beta-synthase Deficiency	1 in 25,000	< 1 in 1,000,000
ypophosphatasia, Autosomal Recessive	1 in 16,000	< 1 in 1,000,000
nclusion Body Myopathy 2	< 1 in 50,000	< 1 in 1,000,000
ovaleric Acidemia	1 in 25,000	< 1 in 1,000,000
oubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
rabbe Disease	1 in 15,000	< 1 in 1,000,000
poamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
aple Syrup Urine Disease Type 1B	1 in 25,000	< 1 in 1,000,000
ledium Chain Acyl-CoA Dehydrogenase Deficiency	1 in 5,900	< 1 in 1,000,000
egalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000	< 1 in 1,000,000
letachromatic Leukodystrophy	1 in 20,000	< 1 in 1,000,000
lucolipidosis IV	< 1 in 50,000	< 1 in 1,000,000
lucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
luscle-eye-brain Disease	< 1 in 12,000	< 1 in 1,000,000
EB-related Nemaline Myopathy	< 1 in 6,700	< 1 in 1,000,000
iemann-Pick Disease Type C	1 in 19,000	< 1 in 1,000,000
iemann-Pick Disease, SMPD1-associated	1 in 25,000	< 1 in 1,000,000
ijmegen Breakage Syndrome	1 in 16,000	< 1 in 1,000,000
orthern Epilepsy	< 1 in 50,000	< 1 in 1,000,000
CDH15-related Disorders		< 1 in 1,000,000
	1 in 5,300	
endred Syndrome	1 in 7,000	< 1 in 1,000,000
EX1-related Zellweger Syndrome Spectrum	1 in 11,000	< 1 in 1,000,000
henylalanine Hydroxylase Deficiency	1 in 5,000	1 in 990,000
KHD1-related Autosomal Recessive Polycystic Kidney Disease	1 in 6,100	< 1 in 1,000,000
olyglandular Autoimmune Syndrome Type 1	1 in 14,000	< 1 in 1,000,000
ompe Disease	1 in 6,300	< 1 in 1,000,000
PT1-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
rimary Carnitine Deficiency	< 1 in 50,000	< 1 in 1,000,000
rimary Hyperoxaluria Type 1	1 in 35,000	< 1 in 1,000,000
rimary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
ROP1-related Combined Pituitary Hormone Deficiency	1 in 11,000	< 1 in 1,000,000
seudocholinesterase Deficiency (Mild Condition)	1 in 2,700	1 in 300,000
ycnodysostosis	< 1 in 50,000	< 1 in 1,000,000
hizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
alla Disease	< 1 in 30,000	< 1 in 1,000,000
egawa Syndrome	< 1 in 50,000	< 1 in 1,000,000
nort Chain Acyl-CoA Dehydrogenase Deficiency	1 in 16,000	< 1 in 1,000,000
ogren-Larsson Syndrome	1 in 9,100	< 1 in 1,000,000
nith-Lemli-Opitz Syndrome	1 in 4,900	1 in 970,000
oinal Muscular Atrophy	Negative for g.27134T>G SNP SMN1: 2 copies 1 in 770	1 in 110,000
eroid-resistant Nephrotic Syndrome	1 in 40,000	< 1 in 1,000,000
ulfate Transporter-related Osteochondrodysplasia	1 in 11,000	< 1 in 1,000,000
PP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
rosinemia Type I	1 in 17,000	< 1 in 1,000,000
sher Syndrome Type 3	< 1 in 50,000	< 1 in 1,000,000
ery Long Chain Acyl-CoA Dehydrogenase Deficiency	1 in 8,800	< 1 in 1,000,000
ary Long Chain Acyrcoa Denyarogenase Denciency	1 111 0,000	~ I III 1,000,000